


WINDOWS TO THE BRAIN

Robin A. Hurley, M.D., L. Anne Hayman, M.D., Katherine H. Taber, Ph.D.
Section Editors

Traumatic Axonal Injury: Atlas of Major Pathways

Katherine H. Taber, Ph.D., Robin A. Hurley, M.D.



Cover and Figure 1. The structures involved in the dorsolateral (pink), anterior cingulate (green), and orbitofrontal (blue) prefrontal-subcortical circuits are color-coded onto the left side of axial magnetic resonance images (A-E). The approximate extent and locations of the classic major long cortical association tracts are color-coded onto the right side of axial magnetic resonance images (A-E) and summarized in cartoon form on a lateral view of the brain (Cover).^{1,2}




Figure 2. Variability (or probability) maps created by transforming the functional anatomy of individual brains into a common anatomic space indicate considerable normal variation.³⁻⁶ On the right is a variability map for primary visual cortex (Brodmann's area 17), with the number of individuals (out of 10) which overlapped. On the left is a probability map (30% threshold) of the corpus callosum, color-coded by cortical area of fiber origin. Note the large areas of overlap, indicating differences in functional anatomy across individuals.

There is increasing evidence that combat-related traumatic brain injuries are a frequent occurrence. Recent studies detailing the most common injuries have found that approximately one-half involved the head or neck.^{9,10} The great majority of injuries were due to explosions. Several studies from the Defense and Veterans Brain Injury Center (DVBIC) of soldiers returning from Afghanistan and Iraq document the occurrence of traumatic brain injury (TBI) in many soldiers.^{11–14} Between January 2003 and February 2005, 59% of returning soldiers treated at Walter Reed Army Medical Center who had been near an explosion while deployed had suffered a traumatic brain injury (44% mild, 56% moderate-severe).¹¹ Common postconcussive symptoms included headache (47%), irritability/aggression (45%), and difficulty with memory (46%) and attention/concentration (41%).¹² A study of 596 active duty soldiers (all serving full-time at regular duty stations in the United States) found that 96 (16.1%) reported an injury while deployed, for which the symptoms (e.g., alternation in or loss of consciousness) were consistent with TBI.¹³ This is similar to an earlier study of active duty soldiers, which found that 13.5% of nonparatroopers reported sustaining a TBI while in the Army.¹⁵ The vast majority of these were mild TBIs, as indicated by either no or only brief loss of consciousness. In most cases, these less severe injuries would not have required medical evacuation.¹⁴ It is well known that civilian mild TBI is under-recognized by both medical personnel and patients, resulting in significant underreporting.¹⁶ There is evidence for a similar situation in the military and concern that combat-related mild TBI may often be unrecognized by both medical personnel and soldiers.^{13,15,17}

Identification of TBI, particularly mild TBI, is often quite challenging. The most common type of injury, and the most likely injury to occur in mild TBI, is traumatic axonal injury (also called diffuse axonal injury).¹⁸ While

magnetic resonance imaging (MRI) is more sensitive than computed tomography (CT) in detecting this type of brain injury, even MRI is often negative.^{18–23} In addition, some areas of injury may become less visible with time.²¹ In such cases, MRI in the subacute and chronic stages is less likely to be positive than if acquired immediately following injury. There is increasing evidence that functional imaging (e.g., cerebral blood flow, cerebral metabolic rate) may be considerably more sensitive to the effects of TBI than structural imaging.^{23–27}

Even small areas of injury within the white matter may have devastating consequences. Knowledge of the locations of major tracts and the brain areas they interconnect is thus critical for understanding clinical symptoms in the context of TBI. White matter tracts of particular importance in neuropsychiatry include those interconnecting areas of cortex (e.g., corpus callosum, association fiber tracts), those connecting areas of cortex to subcortical structures critical for cognitive/emotional functions (e.g., thalamic radiations) and those interconnecting these subcortical areas (e.g., fornix).^{2,28} Tables 1 to 3 summarize the classic functional anatomy of the major white matter pathways important for cognitive and emotional functioning (Figure 1). They are based on recent studies delineating the anatomy of white matter in humans, primarily using diffusion tensor imaging.^{1,2,6–8,29–33}

Intriguing results from sophisticated radioisotope tract-tracing studies in nonhuman primates suggests that there may be significant errors in the classic view of cerebral white matter.^{34,35} For example, this work has delineated three different components (in both location and areas connected) of the superior longitudinal fasciculus. A fourth pathway, which this research group considered to be the arcuate fasciculus, was also identified. A recent diffusion tensor imaging study³⁶ supports the existence of all four of these pathways in humans. Methods for delineating connections within the intact brain are undergoing rapid development and refinement. It is extremely likely that over the next decade, our understanding of the pathways within the brain that are important for cognitive and emotional functioning will change dramatically.

CONCLUSIONS

Care must be taken in applying this summary of functional anatomy to individual patients, as studies comparing pathway topography between subjects have shown considerable normal variability (Figure 2).^{6–8}

Drs. Taber and Hurley are affiliated with the Veterans Affairs Mid Atlantic Mental Illness Research, Education, and Clinical Center, and the Mental Health Service Line, Salisbury Veterans Affairs Medical Center, Salisbury, North Carolina. Dr. Taber is also affiliated with the Division of Biomedical Sciences, Virginia VA School of Osteopathic Medicine, Blacksburg, Virginia, and the Department of Physical Medicine and Rehabilitation, Baylor College of Medicine, Houston, Texas. Dr. Hurley is also affiliated with the Departments of Psychiatry and Radiology, Wake Forest University School of Medicine, Winston-Salem, North Carolina, and the Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas. Address correspondence to Dr. Hurley, Hefner VA Medical Center, 1601 Brenner Avenue, Salisbury, NC 28144; Robin.Hurley@med.va.gov (e-mail).

Copyright © 2007 American Psychiatric Publishing, Inc.

TRAUMATIC AXONAL INJURY

TABLE 1. Commissures Connecting Cortical Areas in Left and Right Hemispheres

Pathway (Areas Connected)	Functional Deficits ²
Corpus Callosum ^{2,7,29,30} Rostrum, genu and anterior body (prefrontal cortex) Mid-body (pre and supplementary motor cortices) Posterior Body (primary motor, primary sensory, parietal cortex) Splenium (temporal and occipital cortices)	Callosal disconnection syndromes Cognitive dysfunctions; alien hand syndromes; apraxias Compulsive/impulsive grasping/reaching due to loss of voluntary inhibition Ideomotor and constructional apraxias; alien hand syndromes; dysomia; agnosias; agraphia; motor neglect; akinesias; tactile-verbal disconnection Visuomotor impairments; optic aphasia; optic ataxia; visual agnosia; memory impairments Minimal information for discrete lesions—left hand apraxia and hearing deficits if injured along with the genu of the corpus callosum
Anterior Commissure ^{2,29,30} Anterior division (olfactory bulbs and nuclei, anterior perforated substance) Posterior division (parahippocampal region, amygdala, inferior temporal and occipital cortex)	

TABLE 2. Long Tracts Connecting Cortical Areas Within Each Hemisphere

Pathway (Areas Connected)	Functional Deficits ²
Superior Longitudinal Fasciculus ^{2,29,30} aka Arcuate Fasciculus Medial long fibers (lateral frontal [Broca's area] to dorsolateral parietal, temporal [Wernicke's area] & occipital cortex) Lateral short fibers (frontal to parietal, parietal to occipital, parietal to temporal cortex)	Left: conduction aphasia; ideational apraxia; depression; anomia Right: left hemispatial neglect
Inferior Longitudinal Fasciculus ^{2,29,30} aka Inferior Occipitotemporal Fasciculus Long fibers (superior, middle and inferior temporal to lingula, cuneus, lateral occipital and posterior occipital cortex) Short fibers (temporal to temporal, occipital to occipital, occipital to parietal cortex)	Disruption of information transfer between visual and limbic/memory areas Left: alexia (if splenium also injured); bilateral tactoverbal dysfunction Left or Right: impaired visual recent memory Bilateral: prosopagnosia Bilateral or Unilateral: visual object agnosia; contralateral visual field hemiachromatopsia Bilateral or Right: visual hypoemotionality
Superior Fronto-Occipital Fasciculus ^{2,29,30} aka Subcallosal Fasciculus, Superior Occipitofrontal Fasciculus (Dorsolateral prefrontal to superior parietal cortex; classic occipital and temporal connections now in question)	Left: akinetic mutism; disordered initiation and preparation of speech movements; transcortical motor aphasia; anomia and reduction in spontaneous speech with normal articulation
Inferior Fronto-Occipital Fasciculus ^{2,29-31} aka Inferior Occipitofrontal Fasciculus (Dorsolateral & ventrolateral prefrontal to posterior temporal and occipital cortex; classic occipital connections now in question)	Seldom injured alone—based on anatomy, injury might cause visuospatial abnormalities; visual recognition abnormalities; topognosia Bilateral: oculomotor apraxia; akinsia Bilateral more than Unilateral: optic ataxia; visual agnosia; impaired visual memory Bilateral or Right: impaired simultaneous perception; impaired spatial relations Right more than Left: impaired orienting of attention—important for retrieval of past information
Uncinate Fasciculus ^{2,29-31} (Orbital and inferior frontal to temporal pole, uncus, hippocampal gyrus and amygdala)	Right: impaired retrieval of episodic (autobiographical) memory Left: impaired retrieval of general knowledge of facts
Cingulum ^{2,29,30} Longest fibers (sub-genu frontal and paraolfactory area to uncus and parahippocampal gyrus) Short fibers (interconnects portions of frontal, parietal and temporal cortex)	Lesion-deficit literature provides no way to distinguish between injury to the cingulum and injury to the cingulate cortex Anterior cingulate cortex (agranular cortex) is motor-related (connections to amygdala; nucleus accumbens; medial dorsal thalamus; dorsolateral prefrontal & parietal cortex); injury may cause lack of emotional affective response to pain, depression, anxiety, akinetic mutism, impaired saccades Posterior cingulate cortex (granular cortex) is sensory-related (connections to temporal, parietal and orbitofrontal cortex); injury may cause Left or Right: Retro-splenial amnesia Right: loss of memory for spatial relationships; topographical disorientation Left: loss of verbal memory; blurring of right sides of objects

This is parallel to the normal variation in size, shape, and location of Brodmann's areas (Figure 2).³⁻⁵ This known phenomenon adds a distinct level of uncertainty in predicting individual functional deficits following a brain injury. It should also be kept in mind that in many places multiple pathways travel close together, making

it likely that a TBI will affect more than one and produce complex symptom clusters. These symptoms may not become evident for extended periods of time. Atlases and other visual external memory aids can assist clinicians in rapid memory recall of functional circuits and areas for review on patient imaging examinations.

TABLE 3. Pathways Connecting Cortical With Subcortical Areas Within Each Hemisphere

Pathway (Areas Connected)	Functional Deficits ²
Internal Capsule ^{2,29,30,34}	
Anterior limb (anterior thalamic connections and frontopontine motor connections)	Anterior limb Bilateral: confusion, impaired initiative; impaired affect; impaired verbal memory
Genu (anterior and inferior thalamic connections and corticonuclear motor connections)	Unilateral: impaired reflexive eye saccades; eye deviation to lesion site
Posterior limb (superior, posterior, inferolateral thalamic connections and corticospinal, corticopontine and corticotegmental motor connections)	Genu Bilateral: somnolence; apathy; amnesia; abulia Unilateral: faciolingual weakness; dysarthria; dysphagia; cognitive impairment; executive dysfunction; contralesional asterixis
Note: Recent studies suggest that connections with frontal cortex extend further posterior in the internal capsule than previously thought, encompassing both the anterior limb and the genu. ^{1,37}	Genu-Posterior limb Unilateral: contralesional motor paresis; dizziness or vertigo Left: verbal memory deficits Posterior limb: Bilateral: pseudobulbar mutism; visual deficits; cortical deafness Unilateral: apathy; impaired consciousness; contralesional hemiparesis/hemiplegia; contralesional anesthesia/ataxia Left: verbal memory deficits Recent memory deficits, with recall more affected than recognition; learning dysfunction Right: nonverbal memory deficits; visual retention disturbances, including deficits in anterograde visual memory, revisualization, visuospatial organization, construction ability, and topographical memory Left: verbal memory deficits; cognitive deficits
Fornix ^{2,29,38}	
Precommissural (hypothalamus, septal nuclei, ventral striatum; orbital and anterior cingulate cortex)	
Postcommissural (anterior nucleus of thalamus, hypothalamus—primarily mamillary body)	

References

- Mori S, Wakana S, Nague-Poetscher LM, et al: MRI Atlas of Human White Matter. New York, Elsevier, 2005
- Aralasmak A, Ulmer JL, Kocak M, et al: Association, commissural, and projection pathways and their functional deficit reported in literature. *J Comput Assist Tomogr* 2006; 30:695-715
- Amunts K, Malikovic A, Mohlberg H, et al: Brodmann's areas 17 and 18 brought into stereotaxic space: where and how variable? *Neuroimage* 2000; 11:66-84
- Amunts K, Weiss PH, Mohlberg H, et al: Analysis of neural mechanisms underlying verbal fluency in cytoarchitectonically defined stereotaxic space—the roles of Brodmann areas 44 and 45. *Neuroimage* 2004; 22:42-56
- Uylings HB, Rajkowska G, Sanz-Arigita E, et al: Consequences of large interindividual variability for human brain atlases: converging macroscopical imaging and microscopical neuroanatomy. *Anat Embryol* 2005; 210:431
- Burgel U, Amunts K, Hoemke L, et al: White matter fiber tracts of the human brain: three-dimensional mapping at microscopic resolution, topography and intersubject variability. *Neuroimage* 2006; 29:1092-1105
- Hofer S, Frahm J: Topography of the human corpus callosum revisited—comprehensive fiber tractography using diffusion tensor MRI. *Neuroimage* 2006; 32:989-994
- Zarei M, Johansen-Berg H, Smith S, et al: Functional anatomy of interhemispheric cortical connections in the human brain. *J Anat* 2006; 209:311-320
- Murray CK, Reynolds JC, Schroeder JM, et al: Spectrum of care provided at an Echelon II medical unit during Operation Iraqi Freedom. *Mil Med* 2005; 170:516-520
- Gondusky JS, Reiter MP: Protecting military convoys in Iraq: an examination of battle injuries sustained by a mechanized battalion during Operation Iraqi Freedom, II. *Mil Med* 2005; 170:546-549
- Okie S: Traumatic brain injury in the war zone. *N Engl J Med* 2005; 352:2043-2047
- Warden DL, Ryan LM, Helmick KM, et al: War neurotrauma: the Defense and Veterans Brain Injury Center (DVBIC) experience at Walter Reed Army Medical Center (WRAMC). *J Neurotrauma* 2005; 22:1178
- Schwab KA, Baker G, Ivins BJ, et al: The Brief Traumatic Brain Injury Screen (BTBIS): investigating the validity of a self-report instrument for detecting traumatic brain injury (TBI) in troops returning from deployment in Afghanistan and Iraq. *Neurology* 2006; 66:A235
- Warden D: Military TBI during the Iraq and Afghanistan wars. *J Head Trauma Rehabil* 2006; 21:398-402
- Ivins BJ, Schwab KA, Warden D, et al: Traumatic brain injury in U.S. Army paratroopers: prevalence and character. *J Trauma* 2003; 55:617-621
- Langlois JA, Rutland-Brown W, Wald MM: The epidemiology

- and impact of traumatic brain injury: a brief overview. *J Head Trauma Rehabil* 2006; 21:375–378
17. Drake AI, McDonald EC, Magnus NE, et al: Utility of Glasgow Coma Scale–Extended in symptom prediction following mild traumatic brain injury. *Brain Inj* 2006; 20:469–475
18. Hurley RA, McGowan JC, Arfanakis K, et al: Traumatic axonal injury: novel insights into evolution and identification. *J Neuropsychiatry Clin Neurosci* 2004; 16:1–7
19. Gutierrez-Cadavid JE: Imaging of head trauma, in *Imaging of the Nervous System*. Edited by Latchaw RE, Kucharczyk J, Moseley ME. Philadelphia, Elsevier Mosby, 2005, pp 869–904
20. Le TH, Gean AD: Imaging of head trauma. *Semin Roentgenol* 2006; 41:177–189
21. Brandstack N, Kurki T, Tenovuo O, et al: MR imaging of head trauma: visibility of contusions and other intraparenchymal injuries in early and late stage. *Brain Inj* 2006; 20:409–416
22. Kurca E, Sivak S, Kucera P: Impaired cognitive functions in mild traumatic brain injury patients with normal and pathologic MRI. *Neuroradiology* 2006; Jun 20 [e-pub ahead of print]
23. Shin YB, Kim SJ, Kin IJ, et al: Voxel-based statistical analysis of cerebral blood flow using technetium-99m ECD brain SPECT in patients with traumatic brain injury: group and individual analyses. *Brain Inj* 2006; 20:661–667
24. Barkai G, Goshen E, Tzila Zwas S, et al: Acetazolamide-enhanced neuroSPECT scan reveals functional impairment after minimal traumatic brain injury not otherwise discernible. *Psychiatry Res* 2004; 132:279–283
25. Gowda NK, Agrawal D, Bal C, et al: Technetium tc-99m ethyl cysteinate dimer brain single-photon emission CT in mild traumatic brain injury: a prospective study. *AJNR Am J Neuroradiol* 2006; 27:447–451
26. Shiga T, Ikoma K, Katoh C, et al: Loss of neuronal integrity: a cause of hypometabolism in patients with traumatic brain injury without MRI abnormality in the chronic stage. *Eur J Nucl Med Mol Imaging* 2006; 33:817–822
27. Pavel D, Jobe T, Devore-Best S, et al: Viewing the functional consequences of traumatic brain injury by using brain SPECT. *Brain Cogn* 2006; 60:211–213
28. Tekin S, Cummings JL: Frontal-subcortical neuronal circuits and clinical neuropsychiatry: an update. *J Psychosom Res* 2002; 53:647–654
29. Catani M, Howard RJ, Pajevic S, et al: Virtual in vivo interactive dissection of white matter fasciculi in the human brain. *Neuroimage* 2002; 17:77–94
30. Jellison BJ, Field AS, Medow J, et al: Diffusion tensor imaging of cerebral white matter: a pictorial review of physics, fiber tract anatomy, and tumor imaging patterns. *AJNR Am J Neuroradiol* 2004; 25:356–369
31. Kier EL, Staib LH, Davis LM, et al: MR imaging of the temporal stem: anatomic dissection tractography of the uncinate fasciculus, inferior occipitofrontal fasciculus, and Meyer’s loop of the optic radiation. *AJNR Am J Neuroradiol* 2004; 25:677–691
32. O’Donnell LJ, Kubicki M, Shenton ME, et al: A method for clustering white matter fiber tracts. *AJNR Am J Neuroradiol* 2006; 27:1032–1036
33. Heiervang E, Behrens TE, Mackay CE, et al: Between session reproducibility and between-subject variability of diffusion MR and tractography measures. *Neuroimage* 2006; 33:867–877
34. Schmahmann JD, Pandya DN: *Fiber Pathways of the Brain*. New York, Oxford University Press, 2006
35. Schmahmann JD, Pandya DN, Wang R, et al: Association fibre pathways of the brain: parallel observations from diffusion spectrum imaging and autoradiography. *Brain* 2007 (e-pub ahead of print)
36. Makris N, Kennedy DN, McInerney S, et al: Segmentation of subcomponents within the superior longitudinal fascicle in humans: a quantitative, in vivo, DT-MRI study. *Cerebral Cortex* 2005; 15:854–869
37. Zarei M, Johansen-Berg H, Jenkinson M, et al: Two-dimensional population map of cortical connections in the human internal capsule. *J Magn Reson Imaging* 2007; 25:48–54
38. Nolte J: *The Human Brain*. St. Louis, Mosby, 2002